

Patient Information Card

Indication for which ▼Clairette (cyproterone acetate / ethinylestradiol) is prescribed:

Clairette is used to treat skin conditions such as acne, very oily skin and excessive hair growth in women of reproductive age. Due to its contraceptive properties it should only be prescribed for you if your doctor considers that treatment with a hormonal contraceptive is appropriate.

You should only take Clairette if your skin condition has not improved after use of other anti- acne treatments, including topical treatments and antibiotics

IMPORTANT INFORMATION ABOUT CLAIRETTE AND RISK OF BLOOD CLOTS

All estrogen-progestagen combination products like Clairette increase the rare but important risk of having a blood clot. The overall risk of a blood clot is small but clots can be serious and may in very rare cases even be fatal.

It is very important that you recognise when you might be at greater risk of a blood clot, what signs and symptoms you need to look out for and what action you need to take.

In which situations is the risk of a blood clot highest ?

- in the first year of using Clairette (including if you are re-starting use after a break of 1 month or more)
- if you are very overweight (body mass index over 30 kg/m²);
- if one of your close relatives (parent or sibling) has had a blood clot at a relatively young age (e.g. below 50)
- if you are older than 35 years
- if you have given birth in the previous few weeks

If you smoke and are over 35 years old you are strongly advised to stop smoking or use a non- hormonal treatment for your acne and/or hirsutism.

Seek urgent medical attention if you experience any of the following:

- Severe pain or swelling in either of your legs that may be accompanied by tenderness, warmth or changes in the skin colour such as turning pale, red or blue. You may be experiencing a **deep vein thrombosis**.
- Sudden unexplained breathlessness or rapid breathing; Severe pain in the chest which may increase with deep breathing; sudden cough without an obvious cause (which may bring up blood); You may be experiencing a serious complication of deep vein thrombosis called a **pulmonary embolism**. This occurs if the blood clot travels from the leg to the lung.
- Chest pain, often acute, but sometimes just discomfort, pressure, heaviness, upper body discomfort radiating to the back, jaw, throat, arm together with a feeling of fullness associated with indigestion or choking, sweating, nausea, vomiting or dizziness. You may be experiencing a **heart attack**.
- Weakness or numbness of the face, arm or leg, especially on one side of the body; sudden confusion; trouble speaking or understanding; sudden loss of vision or blurred vision; severe headache/migraine that is worse than normal. You may be experiencing a stroke.

Watch out for symptoms of a blood clot, especially if you have:

- Just had an operation
- been off your feet for a long time (e.g. because of an injury or illness, or if your leg is in a cast)
- a long journey (e.g. a long-haul flight)

Remember to tell your doctor, nurse or surgeon that you are taking Clairette if you:

- Are due to or have had surgery
- Are asked by a healthcare professional if you are taking any medication

Adverse events

▼Reporting of side effects. This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get.

See www.mhra.gov.uk/yellowcard for how to report side effects.

If you get any side effects, talk to your doctor or nurse. This includes any possible side effects not listed in the package leaflet.

By reporting side effects, you can help provide more information on the safety of this medicine.